

## Applications of the Derivative

### Increasing and Decreasing Functions

**Definition:** A function  $f$  is **increasing on an interval**  $(a,b)$  if  $f(x_2) > f(x_1)$  whenever  $a < x_1 < x_2 < b$ ;  $f$  is **decreasing on an interval**  $(a,b)$  if  $f(x_2) < f(x_1)$  whenever  $a < x_1 < x_2 < b$ .

**Theorem:** a) If  $f'(x) > 0$  on an interval  $(a,b)$ , then  $f$  is increasing on  $(a,b)$ .

b) If  $f'(x) < 0$  on an interval  $(a,b)$ , then  $f$  is decreasing on  $(a,b)$ .

#### **Example 1:** Finding Intervals of Increase and Decrease

Find the intervals where the function  $f(x) = 2 - x^3$  is increasing and those for which it is decreasing.

#### **Example 2:** Finding Intervals of Increase and Decrease

Find the intervals where the function  $f(x) = x^2 - 4x - 12$  is increasing and those for which it is decreasing.

## *Critical Values and Local Extrema*

### **Definition:**

- a) If  $f(c) \leq f(x)$  for all  $x$  in some interval  $(a, b)$ , then  $f(c)$  is a **local** (or relative) **minimum** value for the function  $f$ .
- b) If  $f(c) \geq f(x)$  on an interval  $(a, b)$ , then  $f(c)$  is a **local** (or relative) **maximum** value for the function  $f$ .
- c) The **critical values** (or **critical numbers**) of  $f$  are the values of  $x$  in the domain of  $f$  where  $f'(x) = 0$  or  $f'(x)$  does not exist.

### **Theorem 3: Existence of Local Extrema (Fermat's Theorem)**

If  $f$  is continuous on the interval  $(a, b)$  and  $f(c)$  is a local extremum, then  $c$  is a critical number for the function  $f$ .

**Example 3:** Examine the function  $f(x) = x^3$  at  $x = 0$ . Does the behavior of the graph contradict the theorem on the existence of local extrema?

### **First-Derivative Test for Local Extrema**

Suppose  $c$  is a critical number of a continuous function  $f$ .

- (a) If  $f'$  changes sign from negative to positive at  $x = c$ , then  $f$  has a local minimum at  $c$ .
- (b) If  $f'$  changes sign from positive to negative at  $x = c$ , then  $f$  has a local maximum at  $c$ .
- (c) If  $f'$  does not change sign at  $x = c$ , then  $f$  does not have a local minimum or local maximum at  $c$ .

**Example 4: Finding Maxima and Minima**

For the function  $f(x) = x^3 - 9x^2 + 24x - 10$ , find (a) the critical numbers, (b) local maxima and minima, then (c) sketch the graph.

**Example 5: Finding Maxima and Minima**

For the function  $f(x) = \frac{x^2}{x^2 - 1}$ , find (a) the critical numbers, (b) local maxima and minima, then (c) sketch the graph.

**Example 6: Finding Maxima and Minima**

For the function  $f(x) = e^x - x^2$ , find (a) the critical numbers, (b) local maxima and minima, then (c) sketch the graph.

**Example 7: Finding Maxima and Minima**

For the functions  $f(x) = \sin x + \cos x$ ,  $f(x) = 2x^{1/2} - 4x^{-1/2}$ , and  $f(x) = x^3 - x$ , find (a) the critical numbers, (b) local maxima and minima, then (c) sketch the graph.

Example: Application p106 Ex 5

**Exercise:** For the following functions, find (a) the critical numbers, (b) local maxima and minima, but do not sketch the graph.

a)  $f(x) = x^3$

b)  $f(x) = x^{1/3}$

c)  $f(x) = \frac{2x^2}{x+3}$

d)  $f(x) = \frac{x+2}{x-3}$

e)  $f(x) = 3 - \frac{4}{x} - \frac{2}{x^2}$

f)  $f(x) = x^3(x-5)$

g)  $f(x) = 5x^{1/4} - 2x^{-3/4}$

h)  $f(x) = -\frac{x^3}{x+1}$

## Section 5: The Second Derivative and Graphs

**Definition: The Second Derivative.** If the derivative  $f'$  of a function  $f$  itself has a derivative, we call this the second derivative of  $f$  and use the symbol  $f''$  to denote this. We can also use write the second derivative of  $f$  using Leibniz notation:

$$y'' = \frac{d}{dx} \left( \frac{dy}{dx} \right) = \frac{d^2y}{dx^2}$$

**Notation:** We can denote the second derivative with a variety of notations, including:  $f''(x)$ ,  $\frac{d^2y}{dx^2}$ ,  $y''$ ,  $D_x f'(x)$ ,  $D_x^2 f(x)$  Similarly, we can define higher-order derivatives. The **third derivative of a function**  $y = f(x)$  is denoted

$y''' = f'''(x) = \frac{d}{dx} \left( \frac{d^2y}{dx^2} \right) = \frac{d^3y}{dx^3}$ . In general, the  $n$ th derivative of a function

$y = f(x)$  is denoted  $y^{(n)} = f^{(n)}(x) = \frac{d^n y}{dx^n}$ .

**Exercise:** Find the indicated derivative of the following functions:

1.  $y''$  for  $y = 2x^5 - 6x^4 + 2x - 1$

2.  $f''(x)$  for  $f(x) = \frac{a + bx}{a - bx}$

3.  $f'''(x)$  for  $f(x) = \frac{x}{1 + x}$

## Concavity

Definitions: If  $f'(x)$  is increasing on an interval  $(a,b)$ , then the function  $f$  is said to be **concave upward** on  $(a,b)$ . : If  $f'(x)$  is decreasing on an interval  $(a,b)$ , then the function  $f$  is said to be **concave downward** on  $(a,b)$ .

How do we determine if  $f'$  is increasing or decreasing? By looking at the derivative of the derivative, the second derivative, of course.

**Test for Concavity:** If for each  $x \in (a,b)$ , we have:

- a)  $f''(x) > 0$ , then  $f'(x)$  is increasing on  $(a,b)$  and the graph of  $f$  is concave upward (smiling or bending to the left).
- b)  $f''(x) < 0$ , then  $f'(x)$  is decreasing on  $(a,b)$  and the graph of  $f$  is concave downward (frowning or bending to the right).

**Exercise:** Explain what happens when  $f''(x) = 0$ ?

**Example 1:** The graph of  $f(x) = x^3$  changes concavity from concave downward to concave upward at  $x = 0$ .

**Example 2: Determining Concavity**

Find the intervals where the graph of the function  $f(x) = 2x^3 + 9x^2 - 24x - 10$  is concave upward and those intervals for which the graph is concave downward.

**Example 3: Determining Concavity**

Find the intervals where the graph of the function  $f(x) = x^4 - 8x^2 + 10$  is concave upward and those intervals for which the graph is concave downward.

**Example 4: Determining Concavity**

Find the intervals where the graph of the function  $f(x) = \frac{1-\sqrt{x}}{1+\sqrt{x}}$  is concave upward and those intervals for which the graph is concave downward.

**Exercise: Determining Concavity**

Find the intervals where the graph of the following functions are concave upward and those intervals for which the graph is concave downward.

a)  $f(x) = x^4 - 6x^2 + 1$

b)  $f(x) = \frac{1}{4}x^4 - \frac{1}{2}x^2$

c)  $f(x) = \frac{x+3}{x-3}$

**Inflection Points**

**Definition:** An **inflection point** is a point on the graph of a function where the concavity changes (from upward to downward or downward to upward).  
 $f''$  must change sign at this point.

**Theorem :** If  $(c, f(c))$  is an inflection point of the graph of  $f$ , then either  $f''(c) = 0$  or  $f''$  is undefined at  $x = c$ .

**Exercise:** Find the inflection points of the functions given in Examples 1-4 above.

**Second-Derivative Test for Local Maxima and Minima**

Let  $f$  be a function such that  $f'(c) = 0$  and the second derivative  $f''$  of  $f$  exists on an open interval  $(a, b)$  containing  $c$ .

- a) If  $f''(c) > 0$ , then  $f(c)$  is a relative (local) minimum on  $(a, b)$ .
- b) If  $f''(c) < 0$ , then  $f(c)$  is a relative (local) maximum on  $(a, b)$ .
- c) If  $f''(c) = 0$ , then the test fails (use the first derivative test).

**Example 5: Finding Local Maxima and Minima**

Find any local maxima, local minima for the function  $f(x) = x^2 - \frac{16}{x}$ . Find the inflection points and sketch the graph.

**Example 6: Finding Local Maxima and Minima**

Find any local maxima, local minima for the function  $f(x) = x^{2/3} - 4x^{1/3}$ . Find the inflection points and sketch the graph.

**Exercise:**

- a) Find the intervals where the graph of  $f$  is concave upward, the intervals where the graph of  $f$  is concave downward, and the inflection points for the function  $f(x) = x^4 + 24x^2 + 15x - 12$ .
- b) Find any local maxima, local minima for the function  $f$  defined below. Also find the inflection points. Sketch the graph; include the tangent line at each local extreme point.

$$f(x) = x^3 - 9x^2 + 15x + 10.$$

**Absolute Maxima and Minima**

Definitions:

- a)  $f(c)$  is an **absolute maximum** of  $f$  if  $f(c) \geq f(x)$  for every  $x$  in the domain of  $f$ .
- b)  $f(c)$  is an **absolute minimum** of  $f$  if  $f(c) \leq f(x)$  for every  $x$  in the domain of  $f$ .

**Theorem : Continuous function on a closed interval**

A continuous function  $f$  defined on a closed interval  $[a, b]$  has both an absolute maximum and an absolute minimum value.

### Steps in Finding Absolute Maximum and Minimum Values

- a) Is  $f$  continuous over  $[a, b]$ ?
- b) Find the values of  $x$  for which  $f'(x) = 0$  or is undefined (the critical values).
- c) Evaluate  $f$  at each of the critical values and at the endpoints.
- d) The absolute maximum value of  $f(x)$  on  $[a, b]$  is the largest of the values found in (c).
- e) The absolute minimum value of  $f(x)$  on  $[a, b]$  is the smallest of the values found in (c).

### Second-Derivative Test for Absolute Maximum and Minimum Values

Suppose  $f$  is continuous on an interval  $I$ , and that  $c$  is the only critical value of  $f$  in  $I$ . Then,

- a) if  $f'(c) = 0$  and  $f''(c) > 0$ , then  $f(c)$  is an absolute minimum,
- b) if  $f'(c) = 0$  and  $f''(c) < 0$ , then  $f(c)$  is an absolute maximum,
- c) if  $f'(c) = 0$  and  $f''(c) = 0$ , then the test fails.

### Example 8: Finding Absolute Maximum and Absolute Minimum Values

Find the absolute maximum and absolute minimum for each of the following functions on the specified intervals, if they exist.

- a)  $f(x) = x^2 - 4$  for  $x \in (-\infty, \infty)$ ,  $x \in (-2, 2)$ ,  $x \in [-2, 2]$

b)  $f(x) = \frac{1}{x}$  for  $x \in [-3, 3]$ ,  $x \in [1, 3]$ .